

On inversion formulas and Fibonomial coefficients

A. Krzysztof Kwaśniewski

member of the Institute of Combinatorics and its Applications

Białystok University (*), Faculty of Physics

PL - 15 - 424 Białystok, ul. Lipowa 41, Poland

kwandr@gmail.com

Ewa Krot-Sieniawska

Białystok University, Institute of Computer Science

PL - 15 - 887 Białystok, ul. Sosnowa 64, Poland

ewakrot@wp.pl

(*) former: Warsaw University Division

FECS'08: The 2008 International Conference on Frontiers in Education: WORLDCOMP'08

Summary

A research problem for undergraduates and graduates is being posed as a cap for the prior antecedent regular discrete mathematics exercises. [Here cap is not necessarily CAP=Competitive Access Provider, though nevertheless ...] The object of the cap problem of final interest i.e. array of fibonomial coefficients and the issue of its combinatorial meaning is to be found in A.K.Kwaśniewski's source papers. The cap problem number seven - still opened for students has been placed on Mathmagics page of the first author [<http://ii.uwb.edu.pl/akk/dydaktyka/dyskr/dyskretna.htm>]. The indicatory references are to point at a part of the vast domain of the foundations of computer science in ArXiv affiliation noted as CO.cs.DM. The presentation has been verified in a tutor system of communication with a couple of intelligent students. The result is top secret.Temporarily. [Contact: Wikipedia; Theory of cognitive development].

MCS numbers: 05A19 , 11B39, 15A09

Keywords: inversion formulas, fibonomial coefficients

presented at the Gian-Carlo Rota Polish Seminar

http://ii.uwb.edu.pl/akk/sem/sem_rota.htm

1. In the realm of knownness. Inversion formulas.

Ex.1 Prove that

$$(1) \quad \sum_{k \geq 0} \binom{n}{k} \binom{k}{l} (-1)^{k-l} = \delta_{nl}$$

HINT: For that to do use the following

$$\begin{aligned} (x+1)^n &= \sum_{k \geq 0} \binom{n}{k} x^k \iff x^n = \sum_{k \geq 0} \binom{n}{k} (x-1)^k \iff \\ &\iff x^n = \sum_{k \geq 0} \sum_{l \geq 0} \binom{n}{k} \binom{k}{l} x^l (-1)^{k-l} \iff (1). \end{aligned}$$

Ex.2 Show that

$$(2) \quad \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} \left[\begin{matrix} k \\ l \end{matrix} \right] (-1)^{n-k} = \delta_{nl}, \quad n \geq 0$$

HINT: Use the combinatorial interpretation of Stirling numbers' of the I and the II

kind, $\left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right]$ and $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$, respectively). Also note:

$$x^{\overline{n}} = \sum_{k=0}^n \left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right] x^k, \quad x^n = \sum_{k=0}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} x^{\overline{k}}, \quad x^{\overline{n}} = (-1)^n (-x)^{\overline{n}}.$$

Then

$$x^k = \sum_{l=0}^k \left\{ \begin{smallmatrix} k \\ l \end{smallmatrix} \right\} x^{\overline{l}} \quad \wedge \quad x^{\overline{l}} = (-1)^i (-x)^{\overline{l}}, \quad l \geq 0 \implies (2).$$

Ex.3 Prove that

$$x^n = \sum_{k=0}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} (-1)^{n-k} x^{\overline{k}}, \quad x^{\overline{n}} = \sum_{k=0}^n \left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right] (-1)^{n-k} x^k, \quad n \geq 0.$$

HINT: Use Exercise 2.

Ex.4 $\left(\begin{smallmatrix} n \\ k \end{smallmatrix} \right)_q^{-1} = ?$, for $\left(\begin{smallmatrix} n \\ k \end{smallmatrix} \right)_q = \frac{nq!}{kq!(n-k)q!}$, $nq \equiv \frac{1-q^n}{1-q}$, $n, k \leq 0$.

This problem is solved. Just contact pp.70&106 in [1] (in polish) and note that $\left(\begin{smallmatrix} n \\ k \end{smallmatrix} \right)_q$ counts objects from the $L(n, q)$ lattice. Then one has

$$\left(\left(\begin{smallmatrix} n \\ k \end{smallmatrix} \right)_q \right)^{-1} = \left(\begin{smallmatrix} n \\ k \end{smallmatrix} \right)_q (-1)^{n-k} q^{\binom{n-k}{2}}.$$

Ex.5 Find the numbers $(C_{n,k})^{-1}$ for the sequence $C_{n,k}$ being the unique solution of the recurrence relation, [3]:

$$(3) \quad C_{n+1,k} = C_{n,k-1} + 2C_{n,k} + C_{n,k+1}$$

$$C_{0,0} = 1, \quad C_{k,0} = 0 = C_{n,n+k}, \quad n, k > 0.$$

For $n, k > 0$ one has ([3]): $C_{n,k} = \binom{2n}{n-k} \frac{k}{n}$.

HINT: From the above one has: $C_{0,0}^{-1} = 1$, $C_{k,0}^{-1} = 0$, $C_{n,n+k}^{-1} = 0$ for $n, k > 0$. Then one uses $\sum_{k \geq 0} \binom{2n}{k} \binom{k}{l} (-1)^{k-l} = \delta_{2n,l}$, to get

$$\sum_{s \leq n} \binom{2n}{s} \frac{n-s}{n} \binom{s}{l} \frac{n}{n-s} (-1)^{s-l} = \delta_{nl}, \quad n, l > 0.$$

Ex.6 One can define the Charlier polynomials, as orthogonal polynomial sequence (see [4], the formula 1.13, $a = -1$):

$$P_n(x) = \frac{1}{n!} \sum_{k=0}^n \binom{n}{k} x^{\overline{k}}.$$

Let $x^n = \sum_{k \geq 0} C_{n,k} P_k(x)$. Find the numbers: $C_{n,k}$ and $(C_{n,k})^{-1}$.

HINT: Use Exercise 2 and $x^{\overline{n}} = \sum_{k=0}^n \left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right] (-1)^{n-k} x^k$.

2. Specificity beyond the realm of knownness?

Ex.7 Discover the inversion formula i.e. the array elements $\left(\begin{smallmatrix} n \\ k \end{smallmatrix} \right)_F^{-1}$ for $\left(\begin{smallmatrix} n \\ k \end{smallmatrix} \right)_F$ being the so called fibonomial coefficients, i.e.

$$\left(\begin{smallmatrix} n \\ k \end{smallmatrix} \right)_F = \frac{n_F!}{k_F!(n-k)_F!},$$

for $n_F = F_n$ being the n -th Fibonacci number, ($n, k > 0$). [9,5,10,11,12]
 POSSIBLE SOLUTION: Let us consider the incidence algebra $I(\Pi)$ of the Fibonacci cobweb poset Π and the standard reduced incidence algebra $R(\Pi)$. These were recognized-discovered [Plato's attitude ?] in [5, 6, 7] and investigated there [L.E.J. Brouwer constructivism attitude and constructivism: [http://en.wikipedia.org/wiki/Constructivism_\(learning_theory\)](http://en.wikipedia.org/wiki/Constructivism_(learning_theory))]
 . Let $f : \Pi \times \Pi \rightarrow \mathbb{R}$ be defined as follows

$$f(x, y) = f(k, n) = \begin{cases} \binom{n}{k}_F & k \leq n \\ 0 & k > n \end{cases}$$

for $x, y \in \Pi$, such that the segment $[x, y] = \{z \in \Pi : x \leq z \leq y\}$ is of type (k, n) , i.e. $r(x) = k$, $r(y) = n$. It is obvious that $f \in R(\Pi)$ (and of course $f \in I(\Pi)$). Then for $g = f^{-1}$ being inverse of f in $R(\Pi)$ (also in $I(\Pi)$) one has following [7]

$$(f * g)(k, n) = \sum_{k \leq l \leq n} F_l g(k, l) f(l, n) = \delta_{nk},$$

i.e.

$$(4) \quad \sum_{k \leq l \leq n} F_l \binom{n}{l}_F g(k, l) = \delta_{nk}.$$

Hence in order to discover the magic formula for $\left(\binom{n}{k}_F\right)^{-1}$ one has to find out an explicit formula for $g = f^{-1}$. Right? Maybe it can be recovered using of the standard formula for an inverse element in $I(\Pi)$?, (see for example [1, 8]).

References

- [1] W. Lipski, W. Marek: *Combinatorial Analysis*, **v.59** BM, PWN, Warsaw **1986** (in polish)
- [2] I. Beck: *Partial Orders and the Fibonacci Numbers*, The Fibonacci Quaterly, **26** (1990), pp.272-274.
- [3] L.W. Shapiro: *A Catalan Triangle*, Discrete Math. 14.1 (1976), pp.83-90.
- [4] T.S. Chihara: *An Introduction to Orthogonal Polynomials*, Gordon& Breach, 1978.
- [5] Krot E.: *The first ascent into the Fibonacci Cobweb Poset*, Advanced Studies in Contemporary Mathematics 11 (2005), No. 2, p.179-184, ArXiv: math.CO/0411007, cs.DM <http://arxiv.org/abs/math/0411007>
- [6] Krot-Sieniawska E.: On incidence algebras description of cobweb posets, arXiv:0802.3703, cs.DM <http://arxiv.org/abs/0802.3703>
- [7] E.Krot-Sieniawska: *Reduced Incidence algebras description of cobweb posets and KoDAGs*, ArXiv: 0802.4293, cs.DM <http://arxiv.org/abs/0802.4293>
- [8] Spiegel E., O'Donnell Ch.J.: *Incidence algebras*, Marcel Dekker, Inc. Basel 1997
- [9] Kwaśniewski A.K.: *Towards ψ -extension of Finite Operator Calculus of Rota* Rep. Math. Phys. vol.48 , No3 (2001) pp.304-342 cs.DM cs.NA ArXiv: math/0402078 <http://arxiv.org/abs/math/0402078>

- [10] Kwaśniewski A.K.: *First observations on Prefab posets' Whitney numbers*, Advances in Applied Clifford Algebras Volume 18, Number 1 / February, 2008, p. 57-73. ONLINE FIRST, Springer Link Date, August 10, 2007, arXiv:0802.1696, cs.DM <http://arxiv.org/abs/0802.1696>
- [11] Kwaśniewski A.K.: *On cobweb posets and their combinatorially admissible sequences*, ArXiv:math.Co/0512578v4 21 Oct 2007, submitted to Graphs and Combinatorics; cs.DM <http://arxiv.org/abs/math/0512578>
- [12] Kwaśniewski A.K., Dziemiańczuk M.: *Cobweb posets - Recent Results*, ISRAMA 2007, December 1-17 2007 Kolkata, INDIA, arXiv:0801.3985, cs.DM <http://arxiv.org/abs/0801.3985>